

Remark on the Serre–Swan theorem for non-compact manifolds

G.SARDANASHVILY

Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

E-mail: sard@grav.phys.msu.su

Abstract

The Serre–Swan theorem provides the link between projective modules of finite rank and vector bundles over compact manifolds, and plays a prominent role in non-commutative geometry. Its extension to non-compact manifolds is discussed.

The Serre–Swan theorem had been established for continuous functions and sections [4], but then it was extended to the smooth case (see Proposition 4.1 in Ref. [3] and Proposition 3.1, Theorem 3.2 in Ref. [5]).

Recall that smooth manifolds are assumed to be real, finite-dimensional, Hausdorff, second-countable, and connected. Vector bundles are finite-dimensional.

Theorem 1. (*Serre–Swan*). (i) *The $C^\infty(X)$ -module of global sections of a smooth vector bundle $E \rightarrow X$ over a compact manifold X is a projective module of finite rank.*

(ii) *Any projective $C^\infty(X)$ -module of finite rank is isomorphic to the module of global sections of some smooth vector bundle over X .*

In non-commutative geometry, one therefore thinks of a finite projective $*$ -module over a dense unital $*$ -subalgebra of a C^* -algebra as being a non-commutative vector bundle.

Note that item (i) of Theorem 1 is equivalent to the following well-known theorem (see Theorem 6.5 in Ref. [2] in the continuous case).

Theorem 2. *Let E be a vector bundle over a compact manifold X . There exists a vector bundle $E' \rightarrow X$ such that the Whitney sum $E \oplus E'$ is a trivial vector bundle.*

The key point of the proof of item (i) of Theorem 1 is that any vector bundle over a compact manifold X admits a bundle atlas over a finite covering of X . Proposition IX in Ref. [1] generalizes this assertion to non-compact manifolds as follows.

Proposition 3. *A smooth fibre bundle $Y \rightarrow X$ over an arbitrary manifold X admits a bundle atlas over a finite covering of X .*

Its proof is based on the fact that, for any covering of a manifold X , there exists a refinement $\{U_{ij}\}$, where j and i run through a countable set and a finite set, respectively, such that $U_{ij} \cap U_{ik} = \emptyset$, $j \neq k$. Let $\{U_\xi, \psi_\xi\}$ be a bundle atlas of a fibre bundle $Y \rightarrow X$ over a covering $\{U_\xi\}$ of X . Let $\{U_{ij}\}$ be the above mentioned refinement of this covering and $\{(U_{ij}, \psi_{ij})\}$ the corresponding bundle atlas of $Y \rightarrow X$. Then $Y \rightarrow X$ has the finite bundle atlas

$$U_i \stackrel{\text{def}}{=} \bigcup_j U_{ij}, \quad \psi_i(x) \stackrel{\text{def}}{=} \psi_{ij}(x), \quad x \in U_{ij} \subset U_i. \quad (1)$$

It is readily observed that, if $Y \rightarrow X$ is a vector bundle, the atlas (1) is an atlas of a vector bundle. It follows that every smooth vector bundle admits a finite atlas, and the proof in Refs. [3, 5] of item (i) of Theorem 1 can be generalized straightforwardly to non-compact manifolds. Similarly, the above mentioned Theorem 6.5 in [2] can be extended to non-compact finite-dimensional topological manifolds. The proof of item (ii) in Theorem 1 does not imply the compactness of X .

References

- [1] W.Greub, S.Halperin and R.Vanstone, *Connections, Curvature, and Cohomology*, Vol. 1, (Academic Press, N.Y., 1972).
- [2] M.Karoubi, *K-Theory. An Introduction* (Springer-Verlag, Berlin, 1978).
- [3] G.Landi, An introduction to non-commutative spaces and their geometry, E-print arXiv: hep-th/9701078.
- [4] R.Swan, Vector bundles and projective modules, *Trans. Amer. Math. Soc.* **105** (1962) 264.
- [5] J.Várilly and J.Grasia-Bondia, Connes' noncommutative differential geometry and the standard model, *J. Geom. Phys.* **12** (1993) 223.